

DTIC FILE COPY



AD-A185 019

DEVELOPMENT OF MICROENCAPSULATION
TECHNIQUES

DR. JOHN D. BALDESWIELER

NOVEMBER 26, 1986

U. S. ARMY RESEARCH OFFICE

GRANT NUMBER DAAG29-83-K-0128

CALIFORNIA INSTITUTE OF TECHNOLOGY
PASADENA, CALIFORNIA 91125

DTIC
ELECTE
SEP 23 1987
S D

APPROVED FOR PUBLIC RELEASE;
DISTRIBUTION UNLIMITED.

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

ADA185019

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER ARO 20030.1-CH	2. GOVT ACCESSION NO. N/A	3. RECIPIENT'S CATALOG NUMBER N/A
4. TITLE (and Subtitle) Development of Microencapsulation Techniques		5. TYPE OF REPORT & PERIOD COVERED Final 9/1/83-8/31/86
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Dr. John D. Baldeschwieler		8. CONTRACT OR GRANT NUMBER(s) DAAG29-83-K-0128
9. PERFORMING ORGANIZATION NAME AND ADDRESS California Institute of Technology 1201 E. California Blvd. Pasadena, California 91125		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office Post Office Box 12211 Research Triangle Park, NC 27709		12. REPORT DATE 11/26/86
		13. NUMBER OF PAGES
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) Unclassified
		16a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report) NA		
18. SUPPLEMENTARY NOTES The view, opinions, and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy, or decision, unless so designated by other documentation.		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Phospholipid vesicles Polymerizable phospholipids Spontaneous vesicle formation Liposome		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) When vesicles are formed using phospholipids containing terminal sulphydryl groups it is possible to polymerize the sulphydryl groups by oxidation to form disulfide linkages. Such cross-links lead to a considerable enhancement in vesicle stability for a given size. An important feature of the sulphydryl system is that this type of phospholipid polymer can be readily depolymerized by reducing agents. We have also found formulations which result in spontaneous formation of vesicle structures. Spontaneous vesicle formation reduces many of the difficulties involved in vesicle preparation. Certain combinations of short		

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE(When Data Entered)

20. Abstract(continued)

and long chain lipids give rise to spontaneous vesicle formation. We are currently exploring whether this phenomenon may also occur for short and long chain polymerizable sulfhydryl phospholipids.

Accession For	
NTIS CRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	

QUALITY
INSPECTED
2

UNCLASSIFIED